

Bounded Max-Colorings of Graphs

E. Bampis*

A. Kononov†

G. Lucarelli‡

I. Milis‡

April 10, 2009

Abstract

In a bounded max-coloring of a vertex/edge weighted graph, each color class is of cardinality at most b and of weight equal to the weight of the heaviest vertex/edge in this class. The bounded max-vertex/edge-coloring problems ask for such a coloring minimizing the sum of all color classes' weights.

In this paper we present complexity results and approximation algorithms for those problems on general graphs, bipartite graphs and trees. We first show that both problems are polynomial for trees, when the number of colors is fixed, and H_b approximable for general graphs, when the bound b is fixed. For the bounded max-vertex-coloring problem, we show a $17/11$ -approximation algorithm for bipartite graphs, a PTAS for trees as well as for bipartite graphs when b is fixed. For unit weights, we show that the known $4/3$ lower bound for bipartite graphs is tight by providing a simple $4/3$ approximation algorithm. For the bounded max-edge-coloring problem, we prove approximation factors of $3 - 2/\sqrt{2b}$, for general graphs, $\min\{e, 3 - 2/\sqrt{b}\}$, for bipartite graphs, and 2, for trees. Furthermore, we show that this problem is NP-complete even for trees. This is the first complexity result for max-coloring problems on trees.

1 Introduction

The *bounded max-vertex-coloring* (resp. *bounded max-edge-coloring*) problem takes as input a graph $G = (V, E)$, a weight function $w : V \rightarrow N$ (resp. $w : E \rightarrow N$) and an integer b ; the question of this problem is to find a proper vertex- (resp. edge-) coloring of G , $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$, where each color C_i , $1 \leq i \leq k$, has weight $w_i = \max\{w(u) \mid u \in C_i\}$ (resp. $w_i = \max\{w(e) \mid e \in C_i\}$), cardinality $|C_i| \leq b$, and the sum of colors' weights, $W = \sum_{i=1}^k w_i$, is minimized.

We shall denote the vertex and edge bounded max-coloring problems by $VC(w, b)$ and $EC(w, b)$, respectively. These problems, without the presence of the cardinality bound b , have been already addressed in the literature as *max-vertex-coloring* [20] and *max-edge-coloring* [18] problems; here we denote them by $VC(w)$ and $EC(w)$, respectively. For unit weights we get the *bounded vertex-coloring* [2] and *bounded edge-coloring* [1] problems, denoted by $VC(b)$ and $EC(b)$, respectively. For both unbounded colors cardinalities and unit weights, we have the classical *vertex-coloring* (VC) and *edge-coloring* (EC) problems.

Motivation. Max-coloring problems have been well motivated in the literature. Max-vertex-coloring problems arise in the management of dedicated memories, organized as buffer pools, which is the case for wireless protocol stacks like GPRS or 3G [20, 19]. Max-edge-coloring problems arise in switch based communication systems, like SS/TDMA [4, 17], where messages are to be transmitted through direct connections established by an underlying network. Moreover, max-coloring problems correspond to scheduling jobs with conflicts into a batch scheduling environment [9, 13].

In all applications mentioned above, context-related entities require their service by physical resources for a time interval. However, there exists in practice a natural constraint on the number

*IBISC, CNRS FRE 3190, Université d'Évry, France, bampis@ibisc.univ-evry.fr

†Sobolev Institute of Mathematics, pr Koptiyuga 4, Novosibirsk, Russia, alvenko@math.nsc.ru

‡Dept. of Informatics, Athens University of Economics and Business, Greece, {gluc,milis}@aueb.gr

of entities assigned the same resource or different resources at the same time. Indeed, the number of memory requests assigned the same buffer is determined by strict deadlines on their completion times, while the number of messages and jobs assigned, at the same time, to different channels and machines, respectively, is bounded by the number of the available resources. The existence of such a constraint motivates the study of the bounded max-coloring problems.

Related Work. It is well known that for general graphs it is NP-hard to approximate the VC problem within any constant factor and the EC problem within a factor less than $4/3$; for bipartite graphs both problems become polynomial.

The complexity of the $VC(b)$ problem (known also as Mutual Exclusion Scheduling problem [2]) on special graph classes has been extensively studied (see [14] and the references therein). It is polynomial for trees [16], but NP-complete for bipartite graphs even for three colors [3]. This last result implies also a $4/3$ inapproximability bound for the $VC(b)$ problem on bipartite graphs.

The $VC(w)$ problem is not approximable within a factor less than $8/7$ even for planar bipartite graphs, unless $P=NP$ [9, 19]. This bound is tight for general bipartite graphs, as an $8/7$ -approximation algorithm is also known [7, 19]. On the other hand, the complexity of the problem in trees is an open question, while a PTAS for this case has been presented in [19, 12]. Other results for the $VC(w)$ problem on several graph classes have been also presented in [9, 7, 20, 19, 12, 11].

The $EC(b)$ problem is polynomial for bipartite graphs [4] as well as for general graphs if b is fixed [1]. Moreover, it is implied by the results in [14] that there is a $4/3$ approximation algorithm for the $EC(b)$ problem on general graphs.

The $EC(w)$ problem is not approximable within a factor less than $7/6$ even for cubic planar bipartite graphs with edge weights $w(e) \in \{1, 2, 3\}$, unless $P=NP$ [7]. A simple greedy 2-approximation algorithm for general graphs has been proposed in [17]. The complexity of the $EC(w)$ problem on trees remains also open, while a $3/2$ -approximation algorithm has been recently presented [18].

Known results for the $VC(w, b)$ and $EC(w, b)$ problems have been appeared in the context of batch scheduling for complements of special graph classes (see e.g. [13]). In this context both problems have been shown to be polynomial for general graphs and $b = 2$ [5].

In Table 1 we summarize the best known results for bipartite graphs and trees together with our contribution.

Problem	Bipartite graphs		Trees	
	Lower Bound	Upper bound	Lower Bound	Upper bound
$VC(b)$	$4/3$ [3]	$4/3$	<i>OPT</i> [16]	
$VC(w)$	$8/7$ [9, 7, 19]		open*	<i>PTAS</i> [19, 12]
$VC(w, b)$	$4/3$ [3]	$17/11$	open*	PTAS
$EC(b)$	<i>OPT</i> [4]		<i>OPT</i> [4]	
$EC(w)$	$7/6$ [7]	2 [17]	open*	$3/2$ [18]
$EC(w, b)$	$7/6$ [7]	$\min\{e, 3 - 2/\sqrt{b}, H_b\}^{**}$	NP-complete	2

Table 1: Known and ours (in bold) approximability results for bounded and/or max coloring problems. *Even the complexity of the problem is unknown. **We also show a ratio of $\min\{3 - 2/\sqrt{2b}, H_b\}$ for general graphs. In both cases, the ratio H_b holds only if b is fixed.

Our results and organization of the paper. In this paper we deal with bounded max-coloring problems on general graphs, bipartite graphs and trees. Our interest in bipartite graphs and trees is two-fold. Despite their simplicity, these classes of graphs are important in their own right both from a theoretical point of view but also from the applications' perspective [17, 19].

In the next section, we relate our problems with two well known problems, namely the *list coloring* and the *set cover* problems. We also introduce some useful notation. In Section 3, we deal with the $VC(w, b)$ problem and we give a simple 2-approximation algorithm for bipartite graphs. As a byproduct, we show that this algorithm becomes a $4/3$ -approximation algorithm for

the $VC(b)$ problem, which matches the $4/3$ inapproximability bound. Then, we present a generic scheme that we show to be a $17/11$ -approximation algorithm for bipartite graphs, while it becomes a PTAS for trees as well as for bipartite graphs when b is a fixed constant. In Section 4, we deal with the $EC(w, b)$ problem and we present approximation algorithms of ratios $\min\{3 - 2/\sqrt{2b}, H_b\}$, for general graphs, and $\min\{e, 3 - 2/\sqrt{b}, H_b\}$, for bipartite graphs. More interestingly, we prove that the $EC(w, b)$ problem on trees is NP-complete. Given that the complexity question of $VC(w)$, $EC(w)$ and $VC(w, b)$ problems for trees remains open, this is the first max-coloring problem on trees proven to be NP-hard. Finally, we propose a 2-approximation algorithm for the $EC(w, b)$ problem on trees.

2 Preliminaries and Notation

We first establish a relation between our problems and bounded list coloring.

BOUNDED LIST VERTEX (resp. EDGE) COLORING PROBLEM

INSTANCE: A graph $G = (V, E)$, a set of colors $\mathcal{C} = \{C_1, C_2, \dots, C_k\}$, a list of colors $\phi(u) \subseteq \mathcal{C}$ for each $u \in V$ (resp. $\phi(e) \subseteq \mathcal{C}$ for each $e \in E$), and integers b_i , $1 \leq i \leq k$.

QUESTION: Is there a k -coloring of G such that each vertex u (resp. edge e) is assigned a color in its list $\phi(u)$ (resp. $\phi(e)$) and every color C_i is used at most b_i times?

Clearly, the bounded list coloring problems, denoted by $VC(\phi, b_i)$ and $EC(\phi, b_i)$, are generalizations of the $VC(b)$ and $EC(b)$ problems, as well as of the VC and EC problems, respectively. In the next theorem we summarize some of the known results for the $VC(\phi, b_i)$ and $EC(\phi, b_i)$ problems that we shall use in the rest of the paper.

Theorem 1

- (i) The $VC(\phi, b_i)$ problem is NP-complete even for chains, $|\phi(u)| \leq 2$, for all $u \in V$, and $b_i \leq 5$, $1 \leq i \leq k$ [10].
- (ii) Both $VC(\phi, b_i)$ and $EC(\phi, b_i)$ problems are polynomial for trees if the number of colors k is fixed [15, 8].
- (iii) The $VC(\phi, b_i)$ problem is polynomial for general graphs if $k = 2$ [15].

Using an exhaustive transformation of an instance of the $VC(w, b)$ and $EC(w, b)$ problems to an instance of the $VC(\phi, b)$ and $EC(\phi, b)$ problems (where $b = b_i$, $1 \leq i \leq k$), respectively, and Theorem 1(ii) ([15, 8]) we get next proposition.

Proposition 1 *For a fixed number of colors k , both the $VC(w, b)$ and $EC(w, b)$ problems on trees are polynomial.*

Proof: We give here the proof for the $VC(w, b)$ problem; the proof for the $EC(w, b)$ problem is quite similar. Given a vertex weighted graph $G = (V, E)$ we generate all $\binom{|V|}{k}$ possible combinations for the weights of the k colors. Let $w_1 \geq w_2 \geq \dots \geq w_k$ be the colors' weights in such a combination. For each one of these combinations we construct an instance of the $VC(\phi, b)$ problem on the graph G : is there a k -coloring of the vertices of G such that each color is used at most b times and each vertex $v \in V$ is assigned a color in $\phi(v) = \{C_i : w(v) \leq w_i, 1 \leq i \leq k\}$? A "yes" answer to this instance of the $VC(\phi, b)$ problem corresponds to a feasible solution for the $VC(w, b)$ problem of weight $W = \sum_{i=1}^k w_i$. An optimal solution to the $VC(w, b)$ problem corresponds to the combination where W is minimized.

There are $O(|V|^k)$ combinations of weights to be generated. For a fixed k , by Theorem 1(ii) ([15, 8]), the $VC(\phi, b)$ and $EC(\phi, b)$ problems are polynomial and the proposition follows. ■

Next proposition is based on a relation between $VC(w, b)$ and $EC(w, b)$ problems and the set cover problem.

Proposition 2 *For a fixed bound b , there is an H_b -approximation algorithm for both $\text{VC}(w, b)$ and $\text{EC}(w, b)$ problems on general graphs.*

Proof: In the set cover problem, we are given a universe U of elements, and a collection, $\mathcal{S} = \{S_1, S_2, \dots, S_m\}$, of subsets of U , each one of positive cost c_i , $1 \leq i \leq m$, and we ask for a minimum cost subset of $\mathcal{S}$ that covers all elements of U . For an instance of the $\text{VC}(w, b)$ (resp. $\text{EC}(w, b)$) problem on a graph $G = (V, E)$ we consider the set of vertices V (resp. edges E) corresponding to the universe set U . For each j , $1 \leq j \leq b$, we generate all $\binom{|V|}{j}$ (resp. $\binom{|E|}{j}$) possible subsets of cardinality j of vertices (resp. edges) of G . From all these subsets we get rid those containing adjacent vertices (resp. edges) and we consider the rest corresponding to the set $\mathcal{S}$. For each such subset $S_i \in \mathcal{S}$ we set $c_i = \max\{w(u) | u \in S_i\}$ (resp. $c_i = \max\{w(e) | e \in S_i\}$).

The cardinality of $\mathcal{S}$ is $O(|V|^b)$, since b is $O(|V|)$, and as an H_b -approximation algorithm is known for the set cover problem [6], the proposition follows. ■

Our notation. Given a set S and a positive integer weight $w(s)$ for every element $s \in S$, we denote by $\langle S \rangle = \langle s_1, s_2, \dots, s_{|S|} \rangle$ an ordering of S such that $w(s_1) \geq w(s_2) \geq \dots \geq w(s_{|S|})$. For such an ordering of S and a positive integer b , let $k_S = \lceil \frac{|S|}{b} \rceil$. We define the *ordered b -partition* of S , denoted by $\mathcal{P}_S = \{S_1, S_2, \dots, S_{k_S}\}$, to be the partition of S into k_S subsets, such that $S_i = \{s_j, s_{j+1}, \dots, s_{\min\{j+b-1, |S|\}}\}$, $i = 1, 2, \dots, k$, $j = (i-1)b + 1$. In other words, S_1 contains the b heaviest elements of S , S_2 contains the next b heaviest elements of S and so on; clearly, $S_{k_{|S|}}$ contains the $|S| \bmod b$ lightest elements of S .

By $\text{OPT} = w_1^* + w_2^* + \dots + w_{k^*}^*$ we denote the weight of an optimal solution to the $\text{VC}(w, b)$ or $\text{EC}(w, b)$ problem, where w_i^* , $1 \leq i \leq k^*$, is the weight of the i -th color class. By Δ we denote the maximum degree of a graph.

3 Bounded Max-Vertex-Coloring

In this section we first present a simple 2-approximation algorithm for the $\text{VC}(w, b)$ problem on bipartite graphs. The unweighted variant of this algorithm gives a $\frac{4}{3}$ approximation ratio for the $\text{VC}(b)$ problem on bipartite graphs, which closes the approximability question for this case. Then, we give a generic scheme which becomes a $\frac{17}{11}$ -approximation algorithm for bipartite graphs, a PTAS for bipartite graphs and fixed b , as well as a PTAS for trees. Recall also that by Proposition 2 there is an H_b approximation ratio for general graphs, if b is fixed.

3.1 A simple split algorithm

Let $G = (U \cup V, E)$, $|U \cup V| = n$, be a vertex weighted bipartite graph. Our first algorithm colors the vertices of each class of G separately, by finding the ordered b -partitions of classes U and V . For the minimum number of colors k^* it holds that $k^* \geq \lceil \frac{|U|+|V|}{b} \rceil$ and, therefore, $k = \lceil \frac{|U|}{b} \rceil + \lceil \frac{|V|}{b} \rceil \leq \lceil \frac{|U|+|V|}{b} \rceil + 1 \leq k^* + 1$.

Algorithm SPLIT

1. Let $\mathcal{P}_U = \{U_1, U_2, \dots, U_{k_U}\}$ be the ordered b -partition of U ;
2. Let $\mathcal{P}_V = \{V_1, V_2, \dots, V_{k_V}\}$ be the ordered b -partition of V ;
3. Return the coloring $C = \mathcal{P}_U \cup \mathcal{P}_V$;

Theorem 2 *Algorithm SPLIT returns a solution of weight $W \leq 2 \cdot w_1^* + w_2^* + \dots + w_{k^*}^* \leq 2 \cdot \text{OPT}$ for the $\text{VC}(w, b)$ problem in bipartite graphs.*

Proof: Let $\langle C \rangle = \langle C_1, C_2, \dots, C_k \rangle$ be the colors constructed by Algorithm SPLIT, that is $w_1 \geq w_2 \geq \dots \geq w_k$. Assume, w.l.o.g., that U_x , $1 \leq x \leq k_U$, is the i -th color in $\langle C \rangle$. Let also u be the heaviest vertex of U_x , that is $w(u) = w_i$.

The ordered b -partition of U and V implies that colors $U_1, U_2, \dots, U_{x-1}$ and colors $V_1, V_2, \dots, V_y$, $y = i - x$, appear before color U_x in $\langle C \rangle$. Then, all $(x-1) \cdot b$ vertices of colors $U_1, U_2, \dots, U_{x-1}$

are of weight at least $w(v)$. Also, all $(y-1) \cdot b$ vertices of colors $V_1, V_2, \dots, V_{y-1}$, are of weight at least the weight of the heaviest vertex of color V_y which is at least $w(v)$. Therefore, there are in G at least $(x-1) \cdot b + [(y-1) \cdot b + 1] + 1 = (x+y-2) \cdot b + 2 = (i-2) \cdot b + 2$ vertices of weight at least $w(v) = w_i$. In an optimal solution, these vertices belong into at least $\lceil \frac{(i-2) \cdot b + 2}{b} \rceil = (i-1)$ colors, each one of weight at least w_i . Hence, $w_{i-1}^* \geq w_i$, $2 \leq i \leq k$. Clearly, $w_1 = w_1^*$, since both are equal to the weight of the heaviest vertex of the graph, and as $k \leq k^* + 1$, we obtain

$$\begin{aligned} W &= \sum_{i=1}^k w_i = w_1^* + \sum_{i=2}^k w_i \leq w_1^* + \sum_{i=1}^{k-1} w_i^* \leq w_1^* + \sum_{i=1}^{k^*} w_i^* \\ &= 2 \cdot w_1^* + w_2^* + \dots + w_{k^*}^* \leq 2 \cdot OPT. \end{aligned}$$

■

The complexity of Algorithm SPLIT is dominated by the sorting needed to obtain the ordered b -partitions of U and V in Lines 1 and 2, that is $O(n \cdot \log n)$.

Algorithm SPLIT applies also to the $VC(b)$ problem on bipartite graphs. Moreover, the absence of weights in the $VC(b)$ problem allows a tight analysis with respect to the $\frac{4}{3}$ inapproximability bound.

Theorem 3 *There is a $\frac{4}{3}$ -approximation algorithm for the $VC(b)$ problem on bipartite graphs.*

Proof: Assume, first, that $|U| + |V| \geq 2b + 1$. Then, $k^* \geq \lceil \frac{2b+1}{b} \rceil = 3$ and, since $k \leq k^* + 1$, we get $\frac{k}{k^*} \leq \frac{4}{3}$.

Assume, next, that $b < |U| + |V| \leq 2b$. In this case the optimal solution consists of two or three colors and it is polynomial to decide between them. In fact, it is polynomial to decide if such a bipartite graph can be colored with two colors even for the generalized $VC(\phi, b)$ problem (see also Theorem 1(iii) ([15])).

Assume, finally, that $|U| + |V| \leq b$. Then, an optimal solution consists of either two colors (if $E \neq \emptyset$) or one color (if $E = \emptyset$). ■

3.2 A generic scheme

To obtain our scheme we split a bipartite graph $G = (U \cup V, E)$, $|U \cup V| = n$, into two subgraphs $G_{1,j}$ and $G_{j+1,n}$ induced by the j heaviest and the $n - j$ lightest vertices of G , respectively (by convention, we consider $G_{1,0}$ as an empty subgraph). Our scheme depends on a parameter p such that all the vertices of G of weights $w_1^*, w_2^*, \dots, w_{p-1}^*$ are in a subgraph $G_{1,j}$. This is always possible for some $j \leq b(p-1)$, since each color of an optimal solution for G contains at most b vertices. In fact, for every j , $1 \leq j \leq b(p-1)$, we obtain a solution for the whole graph by concatenating an optimal solution of at most $p-1$ colors for $G_{1,j}$, if there is one, and the solution obtained by Algorithm SPLIT for $G_{j+1,n}$.

Algorithm SCHEME(p)

1. Let $\langle U \cup V \rangle = \langle u_1, u_2, \dots, u_n \rangle$;
2. For $j = 0, 1, \dots, b \cdot (p-1)$ do
3. Split the graph into two vertex induced subgraphs:
 - $G_{1,j}$ induced by vertices $u_1, u_2, \dots, u_j$
 - $G_{j+1,n}$ induced by vertices $u_{j+1}, u_{j+2}, \dots, u_n$
4. If there is a solution for $G_{1,j}$ with at most $p-1$ colors then
5. Find an optimal solution for $G_{1,j}$ with at most $p-1$ colors;
6. Run Algorithm SPLIT for $G_{j+1,n}$;
7. Concatenate the two solutions found in Lines 5 and 6;
8. Return the best solution found;

Lemma 1 *Algorithm SCHEME(p) achieves a $(1 + \frac{1}{H_p})$ approximation ratio for the $VC(w, b)$ problem.*

Proof: Consider the iteration j , $j \leq b \cdot (p-1)$, of the algorithm where the weight of the heaviest vertex in $G_{j+1,n}$ equals to the weight of the i -th color of an optimal solution, i.e. $w(u_{j+1}) = w_i^*$, $1 \leq i \leq p$.

The vertices of $G_{1,j}$ are a subset of those appeared in the $i-1$ heaviest colors of the optimal solution. Thus, an optimal solution for $G_{1,j}$ is of weight $OPT_{1,j} \leq w_1^* + w_2^* + \dots + w_{i-1}^*$.

The vertices of $G_{j+1,n}$ are a superset of those appeared in the $k^* - (i-1)$ lightest colors of the optimal solution. The extra vertices of $G_{j+1,n}$ are of weight at most w_i^* and appear in an optimal solution into at most $i-1$ colors. Thus, an optimal solution for $G_{j+1,n}$ is of weight $OPT_{j+1,n} \leq w_i^* + w_{i+1}^* + \dots + w_{k^*}^* + (i-1) \cdot w_i^* = i \cdot w_i^* + w_{i+1}^* + \dots + w_{k^*}^*$. By Theorem 2, Algorithm SPLIT returns a solution for $G_{j+1,n}$ of weight $W_{j+1,n} \leq (i+1) \cdot w_i^* + w_{i+1}^* + \dots + w_{k^*}^*$.

Therefore, the solution found in this iteration j for the whole graph G is of weight $W_i = OPT_{1,j} + W_{j+1,n} \leq w_1^* + w_2^* + \dots + w_{i-1}^* + (i+1) \cdot w_i^* + w_{i+1}^* + \dots + w_{k^*}^*$.

In all the iterations of the algorithm we obtain p such inequalities for W . By multiplying the i -th, $1 \leq i \leq p$, inequality by $\frac{1}{i \cdot (H_p+1)}$ and adding up all of them, we have $(\sum_{i=1}^p \frac{1}{i \cdot (H_p+1)}) \cdot W \leq OPT$, that is $\frac{W}{OPT} \leq \frac{H_p+1}{H_p} = 1 + \frac{1}{H_p}$. $\blacksquare$

The complexity of the Algorithm SCHEME(p) is $O(bp(f(p) + n \log n))$, where $O(f(p))$ is the complexity of checking for the existence of solutions with at most $p-1$ colors for $G_{1,j}$ and finding an optimal one among them, while $O(n \log n)$ is the complexity of Algorithm SPLIT. Algorithm SCHEME(1) coincides with Algorithm SPLIT. Algorithm SCHEME(2) has simply to check if the $j \leq b$ vertices of $G_{1,j}$ are independent from each other and, therefore, it derives a $\frac{5}{3}$ approximate solution in polynomial time. Algorithm SCHEME(3) has to check and find, a two color solution for $G_{1,j}$, if any. This can be done in polynomial time by Theorem 1(iii) ([15]). Thus, Algorithm SCHEME(3) is a polynomial time $\frac{17}{11}$ -approximation algorithm for the VC(w, b) problem on bipartite graphs.

However, when $p \geq 4$ and b is a part of the instance, finding an optimal solution in $G_{1,j}$ is an NP-hard problem (even for the VC(b) problem [3]). Hence, we consider that b is a fixed constant. In this case, we run an exhaustive algorithm for finding, if any, an optimal solution in $G_{1,j}$ of at most $p-1$ colors. The complexity of such an exhaustive algorithm is $O((p-1)^{b \cdot (p-1)})$ and thus, the complexity of Algorithm SCHEME(p), $p \geq 4$, becomes $O(bp^{bp} + n^2 \log n)$, since bp is $O(n)$. Choosing $\epsilon = \frac{1}{H_p}$, we get $p = O(2^{\frac{1}{\epsilon}})$. Consequently, for fixed b , we have a PTAS for the VC(w, b) problem on bipartite graphs, that is an approximation ratio of $1 + \frac{1}{H_p} = 1 + \epsilon$ within $O(b(2^{\frac{1}{\epsilon}})^{b2^{\frac{1}{\epsilon}}} + n^2 \log n)$ time.

Furthermore, in the particular case of trees, checking the existence of solutions with at most $p-1$ colors for $G_{1,j}$, and finding an optimal one among them, can be done, by Proposition 1, in polynomial time for fixed p . The complexity of our scheme in this case becomes $O(b2^{\frac{1}{\epsilon}}(n^{2^{\frac{1}{\epsilon}}} + n^2 \log n))$. Therefore, the following theorem holds.

Theorem 4 *For the VC(w, b) problem, Algorithm SCHEME(p) is a*
(i) polynomial time $\frac{17}{11}$ -approximation algorithm for bipartite graphs (for $p = 3$),
(ii) PTAS for bipartite graphs if b is fixed,
(iii) PTAS for trees.

4 Bounded Max-Edge-Coloring

In this section we deal with the complexity and approximability of the EC(w, b) problem. We present, first, approximation results for general and bipartite graphs. Then, we prove that the problem is NP-complete for trees and we give a 2-approximation algorithm for this case.

4.1 General and bipartite graphs

We first adapt the greedy 2-approximation algorithm presented in [17] for the EC(w) problem to the EC(w, b) problem.

Algorithm GREEDY

1. **Let** $\langle E \rangle = \langle e_1, e_2, \dots, e_{|E|} \rangle$;
2. **For** $j = 1, 2, \dots, |E|$ **do**
3. **Insert** edge e_j in the first color of cardinality less than b
 which does not contain other edges adjacent to e_j ;

The analysis of Algorithm GREEDY is based on tight bounds on the number of colors in a solution to the $EC(w, b)$ problem.

Proposition 3 *Algorithm GREEDY achieves approximation ratios of $(3 - \frac{2}{\sqrt{2b}})$, on general graphs, and $(3 - \frac{2}{\sqrt{b}})$, on bipartite graphs, for the $EC(w, b)$ problem.*

Proof: We call a solution $\langle C \rangle = \langle C_1, C_2, \dots, C_k \rangle$ to the $EC(w, b)$ problem *nice* if each color C_i , $1 \leq i \leq k$, is of cardinality $|C_i| = b$ or C_i is maximal in the subgraph induced by the edges $E \setminus \bigcup_{j=1}^{i-1} C_j$. We first bound the number of colors in such a solution.

Claim. For the number of colors k in any nice solution to the $EC(w, b)$ problem it holds that:

$$\max\{\Delta, \lceil \frac{|E|}{b} \rceil\} \leq k \leq \begin{cases} \lceil \frac{|E|}{b} \rceil - \lceil \frac{\Delta^2}{2b} \rceil + (2\Delta - 1), & \text{for general graphs} \\ \lceil \frac{|E|}{b} \rceil - \lceil \frac{\Delta^2}{b} \rceil + (2\Delta - 1), & \text{for bipartite graphs} \end{cases}$$

The lower bounds follow trivially. For the upper bounds, let $\langle \mathcal{C} \rangle = \langle C_1, C_2, \dots, C_k \rangle$ be a nice solution, $e = (u, v)$ be an edge in the last color C_k , and E_u and E_v be the sets of edges adjacent to vertices u and v , respectively. By the niceness of the solution $\mathcal{C}$ it follows that edge e does not appear in any color C_i , $1 \leq i \leq k-1$, because $|C_i| = b$ or C_i contains at least one edge in E_u or E_v . Let $W, X, Y \subseteq \{C_1, C_2, \dots, C_{k-1}\}$ such that $W = \{C_i : |C_i| = b\}$, $X = \{C_i : |C_i| < b \text{ and } C_i \text{ contains an edge } e \in E_u\}$ and $Y = \{C_i : |C_i| < b \text{ and } C_i \text{ contains an edge } e \in E_v\}$. Let E_1 be the set of edges in the colors in W and $E_2 = E \setminus E_1$ be the set of edges in the colors in $X \cup Y \cup \{C_k\}$. Then, $k = \frac{|E_1|}{b} + x + y + 1$, where $x = |X|$ and $y = |Y|$.

Assume, w.l.o.g., that $X_1 Y_1 X_2 Y_2 \dots X_l Y_l$ is the order of colors in the nice solution $\langle \mathcal{C} \rangle$, where $X_i \subseteq X$, $Y_i \subseteq Y$, $1 \leq i \leq l$, and X_1 is possibly empty. Let $x_i = |X_i|$ and $y_i = |Y_i|$, $1 \leq i \leq l$.

For general graphs, consider a color $C \in X_i$ and let $S_i = \bigcup_{j=i}^l Y_j$ and $s_i = \sum_{j=i}^l y_j$. The edge $(u, z) \in C \cap E_u$ prevents at most one edge $(v, z) \in S_i \cap E_v$ from being into C . Moreover, each other edge $(p, q) \in C$ prevents at most two edges $(v, p), (v, q) \in S_i \cap E_v$ from being into C . As the colors in S_i contain exactly s_i edges from E_v and all of them are prevented from being into color C , it follows that $|C| \geq \lceil \frac{s_i-1}{2} \rceil + 1$. Therefore, there exist at least $x_i \cdot \lceil \frac{s_i-1}{2} \rceil + x_i$ edges in X_i . In a similar way, by considering a color $C \in Y_i$ there exist at least $y_i \cdot \lceil \frac{t_i-1}{2} \rceil + y_i$ edges in Y_i , where $t_i = \sum_{j=i+1}^l x_j$. Summing up these bounds, and taking into account that $y_l - 1 \geq 0$ (since Y_l is not empty), it follows that

$$\begin{aligned} |E_2| &\geq \sum_{i=1}^l (x_i \cdot \lceil \frac{s_i-1}{2} \rceil + x_i) + \sum_{i=1}^l (y_i \cdot \lceil \frac{t_i-1}{2} \rceil + y_i) \\ &= \frac{x(y-1) - (y_1 + y_2 + \dots + y_{l-1})}{2} + x + y + 1 \\ &\geq \frac{x(y-1) - (y_1 + y_2 + \dots + y_{l-1} + y_l - 1)}{2} + x + y + 1 \\ &= \frac{(x-1)(y-1)}{2} + x + y + 1 \geq \frac{(x+1)(y+1)}{2} + 1 \geq \lceil \frac{(x+1)(y+1)}{2} \rceil. \end{aligned}$$

Therefore, $k = \frac{|E \setminus E_2|}{b} + x + y + 1 \leq \lceil \frac{|E|}{b} \rceil - \lceil \frac{(x+1)(y+1)}{2b} \rceil + x + y + 1$. If $\Delta \leq 2b$ then this quantity is maximized when $x = y = \Delta - 1$ and hence $k \leq \lceil \frac{|E|}{b} \rceil - \lceil \frac{\Delta^2}{2b} \rceil + (2\Delta - 1)$. If $\Delta > 2b$ then the above quantity is maximized when $x = \Delta - 1$ and $y = 0$ and hence $k \leq \lceil \frac{|E|}{b} \rceil - \lceil \frac{\Delta}{2b} \rceil + \Delta \leq \lceil \frac{|E|}{b} \rceil - \lceil \frac{\Delta^2}{2b} \rceil + (2\Delta - 1)$.

For bipartite graphs, the proof is similar. The structure of a bipartite graph allows a tighter bound on the number of edges in the colors in X_i and Y_i . Consider, again, a color $C \in X_i$. For the edge $(u, z) \in C \cap E_u$, there is no edge $(v, z) \in S_i \cap E_v$, while each other edge $(p, q) \in C$ prevents at most one edge (v, p) or (v, q) in $S_i \cap E_v$ from being into C . Thus, $|C| \geq s_i + 1$ and, hence, there exist at least $x_i(s_i + 1)$ edges in X_i . Similarly there exist at least $y_i(t_i + 1)$ edges in Y_i . The rest of the proof is along the same lines, but using these bounds.

We return, now, to the solution, $\langle C \rangle = \langle C_1, C_2, \dots, C_k \rangle$, derived by Algorithm GREEDY. Consider the color C_i and let e_j be the first edge inserted in C_i , i.e. $w_i = w(e_j)$. Let $E_i = \{e_1, e_2, \dots, e_j\}$, G_i be the subgraph of G induced by the edges in E_i , and Δ_i be the maximum degree of G_i .

The solution $\langle C \rangle$ is a nice one, since it is constructed in a First-Fit manner. Moreover, an optimal solution can be also easily transformed into a nice one of the same total weight. For general graphs, by the bounds above, it follows that (i) $i \leq \lceil \frac{|E_i|}{b} \rceil - \lceil \frac{\Delta_i^2}{2b} \rceil + (2\Delta_i - 1)$, and (ii) in an optimal solution the edges of G_i appear in at least $i^* \geq \max\{\Delta_i, \lceil \frac{|E_i|}{b} \rceil\}$ colors, each one of weight at least w_i . Therefore, $\frac{i}{i^*} \leq \frac{\lceil \frac{|E_i|}{b} \rceil - \lceil \frac{\Delta_i^2}{2b} \rceil + (2\Delta_i - 1)}{\max\{\Delta_i, \lceil \frac{|E_i|}{b} \rceil\}}$. By distinguish between $\Delta_i \geq \lceil \frac{|E_i|}{b} \rceil$ and $\Delta_i < \lceil \frac{|E_i|}{b} \rceil$ it follows that in either case $\frac{i}{i^*} \leq 3 - \frac{\Delta_i^2 + 2b}{2b\Delta_i}$. This bound is maximized when $\Delta_i = \sqrt{2b}$, that is $\frac{i}{i^*} \leq 3 - \frac{2}{\sqrt{2b}}$. Thus, $w_i \leq w_{i^*}^* \leq w_{\lceil i/(3 - \frac{2}{\sqrt{2b}}) \rceil}^*$. Summing up these inequalities for all i 's, $1 \leq i \leq k$, we obtain the $(3 - \frac{2}{\sqrt{2b}})$ ratio for general graphs.

A similar analysis yields the $(3 - \frac{2}{\sqrt{b}})$ ratio for bipartite graphs. We present here an example for which the algorithm performs a ratio of exactly $3 - \frac{2}{\sqrt{b}}$ for bipartite graphs. There is, also, an analogous example for general graphs. Consider the bipartite graph shown in Figure 1(a), where $C \gg \epsilon$, and $b = 9$. The weight of the optimal solution shown in Figure 1(b) is $3C + 3\epsilon$. The weight of the solution obtained by Algorithm GREEDY, shown in Figure 1(c), is $7C - \epsilon$. Thus, the ratio for this instance is $\frac{7C - \epsilon}{3C + 3\epsilon} \simeq \frac{7}{3} = 3 - \frac{2}{\sqrt{9}}$.




Figure 1: (a) An instance of the $EC(w, b)$ problem on bipartite graphs, $C \gg \epsilon$, $b = 9$. (b) An optimal solution. (c) The solution obtained by Algorithm GREEDY.

■

Another approximation result for the $EC(w, b)$ problem is obtained by exploiting a general framework, presented in [11], which allows to convert a ρ -approximation algorithm for a coloring problem into an $e \cdot \rho$ -approximation one for the corresponding max-coloring problem, for hereditary classes of graphs. In fact, this framework has been presented for such a conversion from the VC to the $VC(w)$ problem, but it can be easily seen that this applies also for conversions from the EC , $VC(b)$ and $EC(b)$ problems to the $EC(w)$, $VC(w, b)$ and $EC(w, b)$ problems, respectively.

b	General graphs		Bipartite graphs	
3	1.833	H_b	1.833	H_b
4	2.083	H_b	2.000	$3 - 2/\sqrt{b}$
5	2.283	H_b	2.106	$3 - 2/\sqrt{b}$
6	2.423	$3 - 2/\sqrt{2b}$	2.184	$3 - 2/\sqrt{b}$
...	...	$3 - 2/\sqrt{2b}$	...	$3 - 2/\sqrt{b}$
50	2.800	$3 - 2/\sqrt{2b}$	2.717	$3 - 2/\sqrt{b}$
51	2.802	$3 - 2/\sqrt{2b}$	2.718	e
...	...	$3 - 2/\sqrt{2b}$	...	e

Table 2: Approximation ratios for the $EC(w, b)$ problem.

However, this conversion leads to ratios greater than those shown in Table 1 for the $EC(w)$ and $VC(w, b)$ problems. For the $EC(w, b)$ problem on general graphs this approach gives a ratio of at least $\frac{4}{3} \cdot e > 3$, as the EC , and hence the $EC(b)$, problem cannot be approximated within a ratio less than $\frac{4}{3}$. On the other hand, the $EC(w, b)$ problem on bipartite graphs can be approximated, this way, with a ratio of e , as the $EC(b)$ problem is polynomial in this case (see Table 1).

Combining the discussion above with Propositions 3 and 2, it follows that

Theorem 5 *The $EC(w, b)$ problem can be approximated with a ratio of $\min\{3 - 2/\sqrt{2b}, H_b\}$, for general graphs, and $\min\{e, 3 - 2/\sqrt{b}, H_b\}$, for bipartite graphs.*

Note that, the H_b ratio outperforms the other only for $b \leq 5$, for general graphs, and $b = 3$, for bipartite graphs, and, hence, b can be considered as fixed. These ratios are shown in Table 2, for several values of b .

4.2 NP-completeness for trees

We prove first that the bounded list edge-coloring, $EC(\phi, b)$, problem is NP-complete even if the graph $G = (V, E)$ is a set of chains, $|\phi(e)| = 2$, for all $e \in E$, and $b = 5$. We denote this problem as $EC(\text{chains}, |\phi(e)| = 2, b = 5)$.

Proposition 4 *The $EC(\text{chains}, |\phi(e)| = 2, b = 5)$ problem is NP-complete.*

Proof: By Theorem 1(i) ([10]), the $VC(\text{chains}, |\phi(v)| \leq 2, b_i \leq 5)$ problem is NP-complete. Given that the line-graph of a chain is also a chain, it follows that the $EC(\text{chains}, |\phi(e)| \leq 2, b_i \leq 5)$ problem is also NP-complete. The later problem can be easily reduced to the $EC(\text{chains}, |\phi(e)| \leq 2, b = 5)$ problem, where $b_i = b = 5$ for all colors: for every color C_i with $b_i < 5$, add $5 - b_i$ independent edges with just C_i in their lists. This last problem reduces to the $EC(\text{chains}, |\phi(e)| = 2, b = 5)$ problem, where $|\phi(e)| = 2$ for all edges. This can be done by transforming an instance of $EC(\text{chains}, |\phi(v)| \leq 2, b = 5)$ as following: (i) add two new colors C_{k+1} and C_{k+2} , both with cardinality bound $b = 5$, (ii) add color C_{k+1} to the list of every edge e with $|\phi(e)| = 1$, (iii) add ten independent edges and put in their lists both colors C_{k+1} and C_{k+2} . ■

Theorem 6 *The $EC(w, b)$ problem on trees is NP-complete.*

Proof: Our reduction is from $EC(\text{chains}, |\phi(e)| = 2, b = 5)$ problem. We construct an instance of the $EC(w, b)$ problem on a forest $G' = (V', E')$ as follows.

We replace every edge $e = (u, v) \in E$ with a chain of three edges: $e_1 = (u, u')$, $e_2 = (u', v')$ and $e_3 = (v', v)$, where $w(e_1) = w(e_2) = w(e_3) = 1$. Moreover, we create $k - |\phi(e)| = k - 2$ stars of $k - 1$ edges each. We add edges (u', s_t) , $1 \leq t \leq k - 2$, between u' and the central vertex s_t of each of these $k - 2$ stars; thus every star has now exactly k edges. Let $\phi(e) = \{C_i, C_j\}$. The $k - 2$ edges (u', s_t) take different weights in $\{1, 2, \dots, k\} \setminus \{i, j\}$. Let q be the weight taken by an

edge (u', s_t) . The remaining $k - 1$ edges of the star t take different weights in $\{1, 2, \dots, k\} \setminus \{q\}$. In the same way, we add $k - 2$ stars connected to v' . In Figure 2, is shown the u' 's part of this edge-gadget for $e = (u, v)$. For every edge e of G , we add $2(k - 2)$ stars and $2(k - 2)k + 2$ edges.




Figure 2: The gadget for an edge $e = (u, v)$ with $\phi(e) = \{C_i, C_j\}$.

To complete our construction we define f_i to be the frequency of color C_i in the lists of all edges and $F = \max\{f_i | 1 \leq i \leq k\}$. For every color C_i we add $F - f_i$ disconnected copies of the color-gadget shown in Figure 3. Such a gadget consists of an edge $e = (x, y)$ and $k - 1$ stars with $k - 1$ edges each. There are also edges between one of the endpoints of e , say y , and the central vertices of all stars; thus every star has now exactly k edges. The edge e takes weight i and the edges in the stars of such a color-gadget take weights similarly with those in the stars of an edge-gadget. For a color C_i we add $(F - f_i)(k - 1)$ stars and $(F - f_i)(k - 1)k + (F - f_i)$ edges.




Figure 3: A gadget for the color C_i .

The number of stars in the forest G' we have constructed is $2|E|(k - 2) + \sum_{i=1}^k (F - f_i)(k - 1) = k(k - 1)F - 2|E|$, since $\sum_{i=1}^k f_i = 2|E|$. By setting $b' = k(k - 1)F - 2|E| + 5 + F$, we prove that: “There is a k -coloring for $\text{EC}(\phi, b)$ (chains, $|\phi(e)| = 2$, $b = 5$), if and only if, G' has a bounded max-edge-coloring of total weight $\sum_{i=1}^k i$ such that every color is used at most b' times”.

Consider, first, a solution $\mathcal{C}$ to the $\text{EC}(\phi, b)$ problem. We construct a solution $\mathcal{C}'$ for the $\text{EC}(w, b)$ problem as following. Let $e = (u, v) \in E$ be an edge with $\phi(e) = \{C_i, C_j\}$, which, w.l.o.g., appears in the color C_i of $\mathcal{C}$. Put the edges e_1 and e_3 of the edge-gadget for e in color C'_i , while the edge e_2 in color C'_j . After doing this for all edges in E , each color C'_i contains at most $2 \cdot 5 + 1 \cdot (f_i - 5) = f_i + 5$ edges. Next, put the edges with weight i , $1 \leq i \leq k$, from the $k(k - 1)F - 2|E|$ stars into C'_i . Each color C'_i in $\mathcal{C}'$ constructed so far contains at most $k(k - 1)F - 2|E| + f_i + 5 = b' - (F - f_i)$ edges and, by the construction of G' , $\mathcal{C}'$ is a proper coloring. In the $F - f_i$ color-gadgets for C_i there are $F - f_i$ remaining (x, y) edges of weight i , which can still be inserted into color C'_i . Thus, we get a solution for the $\text{EC}(w, b)$ problem of k colors, each one of at most b' edges, and total weight $\sum_{i=1}^k i$.

Conversely, consider a solution $\mathcal{C}'$ to the $\text{EC}(w, b)$ problem. $\mathcal{C}'$ consists of exactly k colors of weights $1, 2, \dots, k$, since each star in G' has k edges and each edge has a different weight in the range $\{1, 2, \dots, k\}$. Thus, all edges of the same weight, say i , should belong in the same color C'_i of $\mathcal{C}'$. Therefore, C'_i contains one edge from each one of the $k(k - 1)F - 2|E|$ stars as well as the $F - f_i$ remaining (x, y) edges of the color-gadgets having weight i . Consider, now, the edges of G' corresponding to the edges e_1 , e_2 and e_3 of the edge-gadget for an edge e with $\phi(e) = \{C_i, C_j\}$.

By the construction of G' and the choice of edge weights, the edges e_1 , e_2 and e_3 should appear into colors C'_i and C'_j . Thus, edges e_1 and e_3 should appear, w.l.o.g., into color C'_i , while e_2 into color C'_j . Therefore, the edge $e \in E$ can be colored by color $C_i \in \phi(e)$. Finally, a color C'_i contains at most 5 edges of type e_1 (or e_3), corresponding to at most 5 edges of E ; otherwise $|C'_i| \geq k(k-1)F - 2|E| + (F - f_i) + (2 \cdot 6 + 1 \cdot (f_i - 6)) > b'$, a contradiction.

To complete our proof for the $EC(w, b)$ problem on trees, let p be the number of trees in G' . We add a set of $p - 1$ edges of weight $\epsilon < 1$ to transform the forest G' into a single tree T . This can be easily done since every tree of G' has at least two vertices. By keeping the same bound b' , it is easy to see that there is a solution for the $EC(w, b)$ problem on G' of weight $\sum_{i=1}^k i$, if and only if, there is a solution for the $EC(w, b)$ problem on T whose weight is equal to $\sum_{i=1}^k i + \lceil \frac{p-1}{b'} \rceil \epsilon$. ■

4.3 A 2-approximation algorithm for trees

In [18] a 2-approximation algorithm for the $EC(w)$ problem on trees has been presented, which is also exploited to derive a ratio of $3/2$ for that problem. This algorithm yields to a solution of Δ colors, $\mathcal{M} = \{M_1, M_2, \dots, M_\Delta\}$. Starting from this solution we obtain a solution to the $EC(w, b)$ problem by finding the ordered b -partition of each color in $\mathcal{M}$. For the sake of completeness we give below the whole algorithm.

Algorithm CONVERT

1. Let T_r be the tree rooted in an arbitrary vertex r ;
2. For each vertex v in pre-order traversal of T_r do
3. Let $\langle E_v \rangle = \langle e_1, e_2, \dots, e_{d(v)} \rangle$ be the edges adjacent to v ,
and (v, p) be the edge from v , $v \neq r$, to its parent;
4. Using ordering $\langle E_v \rangle$, insert each edge in E_v , but (v, p) ,
into the first matching which does not contain an edge in E_v ;
5. Let $\mathcal{M} = \{M_1, M_2, \dots, M_\Delta\}$ be the colors constructed;
6. For $i = 1$ to Δ do
7. Let $\mathcal{P}_{M_i} = \{M_1^i, M_2^i, \dots, M_{k_i}^i\}$ be the ordered b -partition of $\langle M_i \rangle$;
8. Return a solution $\langle \mathcal{C} \rangle = \langle C_1, C_2, \dots, C_k \rangle$, $\mathcal{C} = \bigcup_{i=1}^{\Delta} \mathcal{P}_{M_i}$;

Theorem 7 *Algorithm CONVERT is a 2-approximation one for the $EC(w, b)$ problem on trees.*

Proof: Consider the color C_j in the solution $\langle \mathcal{C} \rangle$ and let e be the heaviest edge in C_j , i.e., $w(e) = w_j$. Let $X \subseteq C_j = \{C_1, C_2, \dots, C_{j-1}\}$ such that each color $C_p \in X$ has (i) $|C_p| = b$, and (ii) all edges of weight at least $w(e)$. Let also $Y = C_j \setminus X$, $|X| = x$ and $|Y| = y$. Clearly, $x + y = j - 1$. Let j^* be the number of colors in an optimal solution of weight at least $w(e)$, that is $w_j = w_{j^*}^*$.

There are at least $x \cdot b + y + 1$ edges of weight at least $w(e)$. These edges in an optimal solution appear in at least $\lceil \frac{x \cdot b + y + 1}{b} \rceil \geq x + 1$ colors, that is, $j^* \geq x + 1$.

We show, next, that all colors in $Y \cup \{C_j\}$ come from $y + 1$ different colors in $\mathcal{M}$. Assume that two of these colors, C_q and C_r , come from the ordered b -partition of the same color $M_t \in \mathcal{M}$. Assume, w.l.o.g., that $w_q \geq w_r$, and let f be the heaviest edge in C_r . Note that C_r may coincide with C_j , while C_q cannot. As $C_q \in Y$, it follows that $|C_q| = b$ and there is an edge $f' \in C_q$ with $w(f') < w(e) \leq w(f)$, a contradiction to the definition of the ordered b -partition of M_t . Therefore, C_j comes from a color $M_i \in \mathcal{M}$, $i \geq y + 1$, that is $e \in M_i$. By the construction of the coloring $\mathcal{M}$, there are at least $i - 1$ edges, adjacent to each other, of weight at least $w(e)$ (i.e., $i - 2$ of them adjacent to e and e itself). These $i - 1$ edges appear in different colors in an optimal solution, that is, $j^* \geq y$.

Combining the two lower bounds for j^* and taking into account that $x + y = j - 1$ we get $j^* \geq \lceil \frac{j}{2} \rceil$. Therefore, $w_j = w_{j^*}^* \leq w_{\lceil \frac{j}{2} \rceil}^*$ and summing up the weights w_j of all colors in $\mathcal{C}$ we get $W = \sum_{j=1}^k w_j \leq 2 \sum_{j=1}^{\lceil k/2 \rceil} w_j^* \leq 2 \sum_{j=1}^{k^*} w_j^* \leq 2OPT$, since $k^* \geq \lceil \frac{k}{2} \rceil$. A tight example for this algorithm, is given in [18], as for large values of b the $EC(w, b)$ coincides with the $EC(w)$ problem.

By a careful analysis, the complexity of both Lines 2 and 6 of the algorithm is $O(n \log n)$, where n is the number of vertices of the tree. ■

References

- [1] N. Alon. A note on the decomposition of graphs into isomorphic matchings. *Acta Mathematica Hungarica*, 42:221–223, 1983.
- [2] B. S. Baker and E. G. Coffman Jr. Mutual exclusion scheduling. *Theoretical Computer Science*, 162:225–243, 1996.
- [3] H. L. Bodlaender and K. Jansen. Restrictions of graph partition problems. Part I. *Theoretical Computer Science*, 148:93–109, 1995.
- [4] G. Bongiovanni, D. Coppersmith, and C. K. Wong. An optimum time slot assignment algorithm for an SS/TDMA system with variable number of transponders. *IEEE Trans. on Communications*, 29:721–726, 1981.
- [5] M. Boudhar and G. Finke. Scheduling on a batch machine with job compatibilities. *Belgian Journal of Oper. Res., Statistics and Computer Science*, 40:69–80, 2000.
- [6] V. Chvátal. A greedy heuristic for the set-covering problem. *Mathematics of Operations Research*, 4:233–235, 1979.
- [7] D. de Werra, M. Demange, B. Escoffier, J. Monnot, and V. Th. Paschos. Weighted coloring on planar, bipartite and split graphs: Complexity and approximation. *Discrete Applied Mathematics*, In Press. (also in Proc. ISAAC’04).
- [8] D. de Werra, A. Hertz, D. Kobler, and N. V. R. Mahadev. Feasible edge coloring of trees with cardinality constraints. *Discrete Mathematics*, 222:61–72, 2000.
- [9] M. Demange, D. de Werra, J. Monnot, and V. Th. Paschos. Time slot scheduling of compatible jobs. *Journal of Scheduling*, 10:111–127, 2007. (also in Proc. WG’02).
- [10] M. Dror, G. Finke, S. Gravier, and W. Kubiak. On the complexity of a restricted list-coloring problem. *Discrete Mathematics*, 195:103–109, 1999.
- [11] L. Epstein and A. Levin. On the max coloring problem. In *5th Workshop on Approximation and Online Algorithms (WAOA’07)*, volume 4927 of *LNCS*, pages 142–155. Springer, 2008.
- [12] B. Escoffier, J. Monnot, and V. Th. Paschos. Weighted coloring: Further complexity and approximability results. *Information Processing Letters*, 97:98–103, 2006. (also in Proc. ICTCS’05).
- [13] G. Finke, V. Jost, M. Queyranne, and A. Sebő. Batch processing with interval graph compatibilities between tasks. *Discrete Applied Mathematics*, 156:556–568, 2008.
- [14] F. Gardi. Mutual exclusion scheduling with interval graphs or related classes. Part II. *Discrete Applied Mathematics*, 156:794–812, 2008.
- [15] S. Gravier, D. Kobler, and W. Kubiak. Complexity of list coloring problems with a fixed total number of colors. *Discrete Applied Mathematics*, 117:65–79, 2002.
- [16] M. Jarvis and B. Zhou. Bounded vertex coloring of trees. *Discrete Mathematics*, 232:145–151, 2001.
- [17] A. Kesselman and K. Kogan. Nonpreemptive scheduling of optical switches. *IEEE Trans. on Communications*, 55:1212–1219, 2007. (also in Proc. GLOBECOM’04).

- [18] G. Lucarelli, I. Milis, and V. Th. Paschos. Maximum edge coloring of trees. *CoRR*, 0901.4002, 2009.
- [19] S. V. Pemmaraju and R. Raman. Approximation algorithms for the max-coloring problem. In *32nd International Colloquium on Automata, Languages and Programming (ICALP'05)*, volume 3580 of *LNCS*, pages 1064–1075. Springer, 2005.
- [20] S. V. Pemmaraju, R. Raman, and K. R. Varadarajan. Buffer minimization using max-coloring. In *15th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA'04)*, pages 562–571, 2004.